



Treatment of Spinal Metastases

INTRABEAM 600 from ZEISS



Seeing beyond

Precise handling in conjunction with palliative treatment – the key benefits of IORT for spinal metastases

For many cancer patients who develop spinal metastases over the course of their disease, percutaneous cement augmentation is a valuable treatment option.¹ By using intraoperative radiation therapy (IORT) during cement augmentation, spinal metastases can be sterilized and the vertebra can be stabilized simultaneously.²

- Using IORT for spinal metastases as an integrated approach enables instant full-weight bearing and restores mobility.^{2,3}
- Long-term stability of the vertebral body due to improved control of local metastases.⁴
- Significant time reduction. Because the radiation is delivered as a single, precisely targeted treatment during surgery, chemotherapy can be started without delay.^{5,6}



1 The tumor is located.



2 Using X-rays, a minimally invasive access point is created.



3 The ZEISS INTRABEAM Needle Applicator is positioned.



4 Radiation (<5 min) of the bone metastasis at its precise location.



5a Stabilization of the vertebra using vertebroplasty or...



5b ... balloon kyphoplasty.



IORT from ZEISS is a new approach for the treatment of spinal metastases. Using this integrated approach, immediate structural stabilization and sterilization of the metastasis can be achieved. Overall, IORT offers benefits for both patient and physician.



The clinical rationale

IORT from ZEISS for irradiation during cement augmentation is an innovative combination of surgery and radiation used to sterilize and stabilize the vertebra simultaneously.⁴ In addition, the high radiation dose with steep dose gradient provides healthy tissue and spinal cord protection. The procedure is a modern treatment option for local tumor control in patients with spinal metastases during palliative treatments. More than 100 operations have been performed using this method.^{7,8,9}

Adapt the radiation to the needs of your patients

To irradiate a tumor cavity, e.g. in the minimally invasive treatment of spinal metastases, ZEISS offers a complete range of applicators in different shapes, sizes and diameters. This versatility enables the physician to exactly adapt the emitted interstitial radiation beam to the form and size of the tumor cavity.



Literature References

- 1 Universitätsmedizin Mannheim. Intraoperative Radiotherapy During Kyphoplasty/Vertebroplasty (Kypho-IORT) Versus EBRT in the Treatment of Painful Vertebral Metastases - a Randomized Prospective Phase III Study -. In: ClinicalTrials.gov [Internet]. Bethesda (MD): National Library of Medicine (US). 2000- [cited 2016 Jul 14]. Available from: <https://clinicaltrials.gov/show/NCT02773966>. NLM Identifier: NCT02773966
- 2 Wenz, F., Schneider, F., Neumaier, C., Kraus-Tiefenbacher, U., Reis, T., Schmidt, R., & Obertacke, U. (2010). Kypho-IORT - a novel approach of intraoperative radiotherapy during kyphoplasty for vertebral metastases. *Radiation Oncology*, 5(1), 11.
- 3 Reis, T., Schneider, F., Welzel, G., Schmidt, R., Bludau, F., Obertacke, U., & Wenz, F. (2012). Intraoperative radiotherapy during kyphoplasty for vertebral metastases (Kypho-IORT): first clinical results. *Tumori*, 98(4), 434–440.
- 4 Schmidt, R., Wenz, F., Reis, T., Janik, K., Bludau, F., & Obertacke, U. (2012). Kyphoplasty and intra-operative radiotherapy, combination of kyphoplasty and intra-operative radiation for spinal metastases: technical feasibility of a novel approach. *International Orthopaedics*, 36(6), 1255–1260.
- 5 Schneider, F., Greineck, F., Clausen, S., Mai, S., Obertacke, U., Reis, T., & Wenz, F. (2011). Development of a Novel Method for Intraoperative Radiotherapy During Kyphoplasty for Spinal Metastases (Kypho-IORT). *International Journal of Radiation Oncology*Biophysics*, 81(4), 1114–1119.
- 6 Miglierini, P., Dam-Hieu, P., & Key, S. (2014). Kypho-IORT: the first French treatment. *Translational Cancer Research*, 3(1).
- 7 Bludau, F. (2014). Kypho-IORT: Intraoperative Radiotherapy in Spinal Metastases: Nine-Month Clinical Results. *The Spine Journal*, 14(11), S163.
- 8 Bludau, F., Reis, T., Schneider, F., Clausen, S., Wenz, F., & Obertacke, U. (2015). Kyphoplastie kombiniert mit intraoperativer Radiotherapie (Kypho-IORT). *Der Radiologe*, 55(10), 859–867.
- 9 Kayser, R., Ender, S. A., Asse, E., Wilhelm, E., Adler, F., Schmidt, R., ... Bludau, F. (2013). Kyphoplastie in Kombination mit intraoperativer Radiotherapie. *Der Orthopäde*, 42(9), 765–771.



Carl Zeiss Meditec AG
Goeschwitzer Strasse 51–52
07745 Jena
Germany
www.zeiss.com/radiotherapy
www.zeiss.com/med/contacts



Carl Zeiss Meditec USA, Inc.
5300 Central Parkway
Dublin, CA 94568
USA
www.zeiss.com/us/radiotherapy
www.zeiss.com/us/med

CAP-en-US_30_020_00041 CZ-IX/2021 Printed in the United States. United States edition. Only for sale in selected countries. The contents of the brochure may differ from the current status of approval of the product or service offering in your country. Please contact our regional representatives for more information. Subject to change in design and scope of delivery and as a result of ongoing technical development. INTRABEAM is either a trademark or registered trademark of Carl Zeiss Meditec AG or other companies of the ZEISS Group in Germany and/or other countries.
© Carl Zeiss Meditec USA, Inc., 2021. All rights reserved.